



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

ing strychnine) has been found very useful in killing the fishes of these pools, often not to be captured in any other way. Mr. Greeley found a good substitute for this poison in the commercial chloride of lime.

D. S. J.

**Development of Brain Structures in *Amia*.** — A. C. Eyclesheimer and B. M. Davis give in the *Journal of Comparative Neurology* a valuable study of "The Early Development of the Epiphysis and Paraphysis in *Amia*." The paper indicates that much is still to be known as to the origin of epiphysial outgrowths from brain structures.

D. S. J.

**Scapanorhynchus and Mitsukurina.** — In the *Annals and Magazine of Natural History*, Mr. A. S. Woodward, of the British Museum, has a note on *Mitsukurina owstoni* Jordan, an extraordinary lamnoid shark with a long flat blade on its snout, lately described from the deep waters of Japan.

Mr. Woodward shows that *Mitsukurina* is very closely related to the Cretaceous genus *Scapanorhynchus*, of which species are known from Mount Lebanon and from the chalk of England.

*Mitsukurina* and *Scapanorhynchus* agree in the elongate, blade-like snout, which is, however, longest in *Scapanorhynchus*. The skeleton, dentition, and gill openings seem to be similar in the two genera, and there appear to be no great differences in the fins. The dense shagreen is also similar in the two; the structure of the basal cartilages of the fins in *Scapanorhynchus* is unknown; nor is anything known of the claspers.

Mr. Woodward concludes that *Mitsukurina* is probably identical with his genus, *Scapanorhynchus*, this name being of prior date. On the other hand, it may be urged that this identity is not proved, and that the specific differences are considerable. There are great disadvantages in the identification of recent fishes with fossil genera which are more or less imperfectly known.

More complete knowledge of the extinct forms often shows that the recent species have undergone such differentiation as should constitute generic difference. I think it, therefore, better to retain for the recent shark the name *Mitsukurina*, although recognizing its close relationship to its Cretaceous homologue.

The family *Mitsukurinidæ* is supposed to differ from *Carchariidæ* (*Odontaspididæ*) in the presence of a *Polyodon*-like snout, and perhaps in the structure of its fins and claspers. The writer knows too